

Transport anomaly in the low energy regime of spin chains

Keiji Saito

*Department of Applied Physics, School of Engineering
University of Tokyo, Bunkyo-ku, Tokyo 113-8656, Japan
()*

The anomalous thermal conductivity in spin chains observed in experiments is studied for the low temperature regime. In the effective dynamics with most realistic perturbations, the so-called Umklapp terms is irrelevant to reduce mean free path in the energy transport at even finite temperatures. This is consistent with large conductivities found in recent experiments. The Drude weight which is the prefactor in the divergent conductivity is calculated, and the temperature dependence is discussed.

PACS numbers: 75.10.Jm, 75.40.Gb

I. INTRODUCTION

Recent experiments on transport phenomena in low dimensional spin systems have attracted much interest^{1,2,3,4,5,6,7,8,9}. There the behavior of conductivity is sensitively affected by the magnetic phases. Therefore the thermal conductivity can be a strong experimental instrument to investigate the magnetic properties in materials. In experiments, magnetic energy transport is much larger than the phononic transport at most temperatures. This fact is considered to be attributed to large exchange coupling constants and nondiffusive behavior of transport. Sr_2CuO_3 described by the isotropic Heisenberg chain has large exchange coupling constant $J \sim 2000\text{K}$ and the magnon can transmit energy even at the order of 100K . The mean free path is experimentally estimated in the similar way as the Peierls-Boltzmann approach for phononic transport^{10,11}, and nondiffusive behavior is found. In another isotropic Heisenberg chain CuGeO_3 , where the mean free path is calculated as 500 times lattice size. It is noteworthy that even when the alternation between bonds occurs, the magnetic transport may be still dominant and nontrivial enhancement of conductivity is observed^{3,15}. Spin-ladder system $(\text{Sr}, \text{La}, \text{Ca})_{14}\text{Cu}_{24}\text{O}_{41}$ and $\text{La}_5\text{Ca}_9\text{Cu}_{24}\text{O}_{41}$ also show extremely large mean free path^{4,5,6,7}.

As studied in classical systems, an integrable system shows ballistic energy transport¹². Even if the energy current is not conserved, the low-dimensional systems with conserved quantities tend to show the divergence of conductivity¹³. These tendencies are also valid in quantum systems^{14,15,16,17,18,19,20,21}. The isotropic Heisenberg chain is the integrable system and exactly shows the diverging conductivity as proved by Zotos et al.²¹. This divergence is a direct result of conserved energy current. However in most materials in experiments, the perturbations like a bond-alternation and a next nearest neighbor interaction exist, so that the energy current is no more conserved. Nevertheless the experiments indicate divergent conductivities. Motivated by these experimental facts, Alvarez and Gros investigated whether the experimentally indicated divergent conductivity really occurs or not. They numerically found nonvanishing Drude

weight at finite low temperatures for the alternate spin chain with frustration and the ladder system¹⁶, although it may vanish at very high temperatures in the thermodynamic limit¹⁸. Numerical finding of finite Drude weight indicates the existence of divergent conductivity at low temperature regime. Stimulated by these numerical studies, in this paper we study the possible mechanism to explain nondiffusive behavior in the presence of perturbations to the Heisenberg chain. We focus on the low energy regime²² by employing the effective dynamics as adopted for the Heisenberg chain by Meisner et al.¹⁸. As a result, we find the effective energy current becomes conserved in spite of the existence of so-called Umklapp terms so that the conductivity diverges. This ballistic nature is a common characteristic for most realistic perturbations as far as the low energy regime is considered. This is a remarkable aspect in the sense that the perturbations is irrelevant to decrease the mean free path although they can be relevant to cause various magnetic phases in the equilibrium state. This is very important to explain the anomalous large conductivities of one-dimensional magnetic systems. We further calculated the quantitative value of the Drude weight of conductivity using the path integral approach, and found it to be finite.

II. EFFECTIVE ENERGY CURRENT IN LOW TEMPERATURE REGIME

We first consider an alternate spin chain whose Hamiltonian is

$$\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_1, \quad (1)$$

$$\mathcal{H}_0 = J \sum_{\ell} \mathbf{S}_{\ell} \cdot \mathbf{S}_{\ell+1}, \quad \mathcal{H}_1 = J\delta \sum_{\ell} (-1)^{\ell+1} \mathbf{S}_{\ell} \cdot \mathbf{S}_{\ell+1}. \quad (2)$$

The total Hamiltonian is divided into the isotropic part $\mathcal{H}_0$ and the dimerized part $\mathcal{H}_1$. Thermal conductivity $\kappa(T)$ at a temperature $T (= 1/\beta)$ is calculated by the Green-Kubo formula which reads as²³,

$$\kappa(T) = \lim_{\omega \rightarrow 0} \kappa(\omega) = \frac{\beta}{2L} \int_{-\infty}^{\infty} dt e^{-i\omega t} \int_0^{\beta} d\tau \langle j(t - i\tau) j \rangle, \quad (3)$$

where $j(\dots)$ is the total current operator in the Heisenberg picture. From the continuity equation of energy, the total current operator j is written as $j = -i \sum_{\ell} [h_{\ell}, h_{\ell+1}]$ with the local Hamiltonian $h_{\ell} := J(1 + (-1)^{\ell+1} \delta) \mathbf{S}_{\ell} \cdot \mathbf{S}_{\ell+1}$ ¹⁵. In the absence of the alternation, j is the conserved quantity²¹, whereas the existence of alternation does not allow the conservation, i.e.,

$$[j_{(\delta=0)}, \mathcal{H}_0] = 0, \text{ but } [j, \mathcal{H}] \neq 0. \quad (4)$$

At low energy regime, the Hamiltonian can be effectively represented by the boson field^{25,26},

$$\mathcal{H}_B = \int dx \left[\pi v p^2 + \frac{v}{4\pi} (\partial_x \Theta_+(x))^2 - B \cos \Theta_+(x) \right], \quad (5)$$

where $v = \frac{J\pi}{2}$ and $B = J\delta$. The Boson field is defined using the fermion field Ψ_R and Ψ_L and satisfies the commutation relation as,

$$\frac{1}{2\pi} \frac{\partial \Theta_{\pm}}{\partial x} = : \Psi_R^{\dagger}(x) \Psi_R(x) : \pm : \Psi_L^{\dagger}(x) \Psi_L(x) :, \quad (6)$$

$$p(x) = -\frac{1}{4\pi} \partial_x \Theta_-(x), \quad (7)$$

$$[\Theta_+(x), \Theta_-(x')] = 2\pi i \operatorname{sgn}(x - x'). \quad (8)$$

We define the local Hamiltonian density at x as, $h(x) = \pi v p^2 + \frac{v}{4\pi} \left(\frac{\partial \Theta_{\pm}}{\partial x} \right)^2 - B \cos \Theta_+(x)$. Using the relations,

$$\frac{\partial \Theta_+}{\partial t} = -2\pi v p, \quad \frac{\partial p}{\partial t} = -\frac{v}{2\pi} \partial_{x^2} \Theta_+(x) + B \sin \Theta_+, \quad (9)$$

we find the Heisenberg equation of motion given by,

$$\frac{\partial h(x, t)}{\partial t} = -\frac{v^2}{2} \partial_x (p \partial_x \Theta_+ + (\partial_x \Theta_+) p). \quad (10)$$

This leads the effective total energy current written as,

$$j_B = v^2 \int dx p(x) \partial_x \Theta_+(x). \quad (11)$$

Interestingly, this effective current operator j_B does not include the perturbation term (Umklapp term) $B \cos \Theta_+(x)$. Furthermore we find that even if the energy current is not rigorously conserved (4), it behaves as the conserved quantity in the low temperature regime, i.e.,

$$[j, \mathcal{H}] \neq 0, \text{ but } [j_B, \mathcal{H}_B] = 0. \quad (12)$$

The perturbation term $B \cos \Theta_+(x)$ causes the Umklapp process in the scattering between fermions²⁷. Nevertheless at the level of the energy transport, this process is irrelevant to decrease the mean free path at finite low temperatures.

This situation is also the case in the other realistic perturbations. We consider the two following examples²⁸,

$$\mathcal{H}_2 = J\alpha \sum_{\ell} \mathbf{S}_{\ell} \cdot \mathbf{S}_{\ell+2}, \quad \mathcal{H}_3 = J\gamma \sum_{\ell} (-1)^{\ell} S_{\ell}^z, \quad (13)$$

The perturbation $\mathcal{H}_2$ causes the frustration and the non-trivial phases appears as a function of α ²⁸. Recent numerical studies treat this perturbation and the effect of frustration is investigated^{16,18}. $\mathcal{H}_3$ is the term of staggered magnetic field. The effective expressions at the low energy regime for the perturbations (13) are written as,

$$\mathcal{H}_2 \sim \int dx c_2 (\partial_x \Theta_+)^2 + c'_2 \cos(2\Theta_+), \quad (14)$$

$$\mathcal{H}_3 \sim c_3 \int dx \cos(2\Theta_+), \quad (15)$$

with constants, c_2 , c'_2 and c_3 . These cases have the same form as the Hamiltonian (5), and the effective total current is easily proved to be conserved. This feature is generally common for the Hamiltonian with the form,

$$\int dx \left[a p^2 + b (\partial_x \Theta_+(x))^2 + \sum_{\ell} c_{\ell} \cos(d_{\ell} \Theta_+(x) + \phi_{\ell}) \right]. \quad (16)$$

Here a, b, c_{ℓ}, d_{ℓ} , and ϕ_{ℓ} are constants. Thus we conclude that the effective energy current is conserved in most realistic spin chains at least as far as we consider in the low temperature regime.

III. DRUDE WEIGHT

Now let us calculate the prefactor of thermal conductivity, i.e., the Drude weight D_{th} in the low energy regime and confirm that finite value of D_{th} really exists. Since the energy current does not depend on a time, we write the Green-Kubo formula as,

$$\kappa = D_{\text{th}} \delta(\omega) \quad (17)$$

$$D_{\text{th}} = \beta\pi/L \int_0^{\beta} d\tau \int_0^L dx \int_0^L dx' \langle j_B(x, \tau) j_B(x', 0) \rangle, \quad (18)$$

where $j(x, \tau)$ is the local energy current at x and the imaginary time τ in the Heisenberg picture. The current is expressed as $j(x, \tau) = \frac{-iv}{2\pi} \partial_x \Theta_+(x, \tau) \partial_{\tau} \Theta_+(x, \tau)$. We confine ourselves in the alternate spin chain effectively described by Hamiltonian (5). At very low temperatures, the Hamiltonian (5) is approximated by the mean field theory³⁰;

$$\cos(\Theta_+) \sim e^{-(\Theta_+^2)/2} \left(1 - \frac{\Theta_+^2 - \langle \Theta_+^2 \rangle}{2} \right). \quad (19)$$

The Hamiltonian is reduced to,

$$\mathcal{H}_B \sim \int dx \left[\frac{v}{4\pi} (\partial_x \Theta_+(x))^2 + C \Theta_+^2 + \pi v p^2 + \text{const.} \right], \quad (20)$$

where $C = B e^{-(\Theta_+^2)/2}$ and $\langle \Theta_+^2 \rangle$ is the average value at the ground state which is self-consistently determined.

This approximation yields the accurate ground state energy within 10 percent error in comparison with the exact numerical data²⁶. Thus we expect that this approximation well describes properties in very low temperatures. Using the Hamiltonian (20), we calculate the Drude weight D_{th} using the path integral (e.g.,²⁹). The Drude weight is given by explicit calculation of

$$D_{\text{th}} = \frac{-v^2}{4\pi\beta L^3} \int \int \int d\tau dx dx' \partial_{x_1} \partial_{x_3} \partial_{\tau_2} \partial_{\tau_4} \sum_{n_1, \dots, n_4} \sum_{k_1, \dots, k_4} e^{i(k_1 x_1 + k_2 x_2 + k_3 x_3 + k_4 x_4)} e^{i(\omega_{n_1} \tau_1 + \omega_{n_2} \tau_2 + \omega_{n_3} \tau_3 + \omega_{n_4} \tau_4)} \times \langle \Theta_+(k_1, n_1) \Theta_+(k_2, n_2) \Theta_+(k_3, n_3) \Theta_+(k_4, n_4) \rangle, \quad (21)$$

where we take $x_1 = x_2 = x, x_3 = x_4 = x', \tau_1 = \tau_2 = \tau$, and $\tau_3 = \tau_4 = 0$. k_m and ω_n are the wave number and the Matsubara frequency defined as $k_m = \frac{2\pi m}{L}$ and $\omega_n = \frac{2\pi n}{\beta}$, respectively. $\Theta_+(k_m, n)$ is the Fourier transformation at the mode k_m and ω_n . The four point correlation is calculated by the Wick's theorem using the two point correlation $\langle |\Theta_+(k_m, n)|^2 \rangle = \frac{2\pi v}{\omega_n^2 + v^2 k_m^2 + 4\pi v C}$. We next carry out the exact summation over n_p ($p = 1, \dots, 4$) using technique of the complex integral. We finally obtain the simple expression of the Drude weight with $a = \frac{\beta}{2\pi} \sqrt{v^2 k^2 + 4\pi v C}$,

$$D_{\text{th}} = \frac{v^4 \beta^2}{8} \int_{-\infty}^{\infty} dk k^2 \left[\left(\frac{1}{e^{2\pi a} - 1} + \frac{1}{e^{-2\pi a} - 1} \right)^2 - \left(\frac{1}{e^{2\pi a} - 1} - \frac{1}{e^{-2\pi a} - 1} \right)^2 \right] \\ = 2vT \int_0^{\infty} dx \frac{x^2}{\sinh^2 \sqrt{x^2 + \pi \beta^2 v C}}. \quad (22)$$

In the case of isotropic Heisenberg chain $C = 0$, $D_{\text{th}} = \frac{v\pi^2 T}{3}$ because of $\int_0^{\infty} dx \frac{x^2}{\sinh^2 x} = \frac{\pi^2}{6}$, which completely agrees with the asymptotic behavior of exact calculation by the Bethe ansatz²⁴. This result is also given by the conformal transformation^{18,31}. In the presence of finite alternation δ , however, the Hamiltonian does not have conformal symmetry.

Eq. (22) shows that the Drude weight is non-vanishing for finite alternation δ . This is quite crucial because the existence of D_{th} assures the anomalous energy transport. The energy gap ΔE is roughly estimated as $\Delta E \sim 2\sqrt{\pi v C}$. Thus for very low temperature regime $T \ll \Delta E$, eq.(22) shows exponential decrease with power law prefactor due to the finite gap

$$D_{\text{th}} \sim \frac{v\beta^2}{32} \int_{-\infty}^{\infty} dp p^2 e^{-\beta \left(\Delta E + \frac{p^2}{2\Delta E} \right)} \\ = \frac{v}{32} \sqrt{\frac{\pi \Delta E^3}{T}} e^{-\beta \Delta E} \quad T \ll \Delta E. \quad (23)$$

Exponential decrease is simply due to exponential decrease of thermal excitation. The power law prefactor

$T^{-1/2}$ is also seen in the electric and spin conductivity in one-dimensional gapped systems^{32,33}. This temperature dependence will be the characteristic in transport phenomena in the low-dimensional gapped systems.

IV. SUMMARY

We find the possible mechanism of anomalous transport in low energy region of realistic spin chains. In the isotropic Heisenberg chain, the energy current is exactly conserved so that the Green-Kubo formula trivially diverges. On the other hand, the presence of perturbations, the Umklapp terms appear. However this term is irrelevant to the energy current of long wave length at low temperatures. We considered the quantitative estimation of the Drude weight for the alternate chain, and find exponential decrease of Drude weight at low temperature. This aspect is consistent with the numerical studies^{16,18}.

Here we should make clear the difference between classical nonintegrable Fermi-Pasta-Ulam (FPU) chains¹³ where the mechanism of anomalous transport is still argued. In this classical system, anomalous heat transport is attributed to a slow relaxation due to the total momentum conservation, whereas the auto-correlation function of energy current vanishes in the long time limit, i.e., the mixing property is satisfied. However, in perturbed quantum Heisenberg spin chains we discuss here, Numerical evidence of the finite Drude weight¹⁶ means the violation of the mixing property. Thus the mechanism in quantum spin chains is different from the one in classical FPU system.

The alternate spin chain directly related to the spin-Peierls system. In the realistic mechanism, the alternation is formed by the fluctuation of the phonon. In this sense, in more realistic treatment, δ must be exchanged by the time-dependent bond length between atoms. In the case where the bond fluctuation is not negligible, how the magnetic energy transport is affected is also interesting.

note added– After submitting this paper, we find the preprint³⁴ where the Drude weight in the dimerized xy spin chain and two-leg ladder system are extensively studied. The paper includes the similar formula as (22) in these systems.

ACKNOWLEDGMENTS

The author would like to thank S. Miyashita and X. Zotos for useful comments. He also expresses thanks the referees for suggestions to interpret the formula (22). The present work is supported by Grand-in-Aid for Scientific Research from Ministry of Education, Culture, Sports, Science, and Technology of Japan.

- ¹ A.V.Sologubenko, K. Gioannò, H. R. Ott, A. Vietkine, and A. Revcolevschi, Phys. Rev. B **62** R6108 (2000), Phys. Rev. B **64** 054412-1 (2001).
- ² Y. Ando, J. Takeya, D. L. Sisson, S. G. Doettinger, I. Tanaka, R. S. Feigelson, and A. Kapitulnik, Phys. Rev. B **58** R2913 (1998).
- ³ M. Hofmann, T. Lorenz, A. Freimuth, G. S. Uhrig, H. Kageyama, Y. Ueda, G. Dhalenne and A. Revcolevschi, Physica B **312-313** 597 (2002).
- ⁴ K. Kudo, S. Ishikawa, T. Noji, T. Adachi, Y. Koike, K. Maki, S. Tsuji, and K. Kumagai, J. Phys. Soc. Jpn **70** 437 (2001).
- ⁵ A.V.Sologubenko, K. Gioannò, H. R. Ott, U. Ammerahl, A. Revcolevschi, D. F. Brewer, and A. L. Thomson, Physica B **284-288** 1595 (2000).
- ⁶ A.V.Sologubenko, K. Gioannò, H. R. Ott, U. Ammerahl, and A. Revcolevschi: Phys. Rev. Lett. **84** 2714 (2000).
- ⁷ C. Hess, C. Baumann, U. Ammeral, B. Büchner, F. Heidrich-Meisner, W. Brenig, and A. Revcholevschi, Phys. Rev. B **64** 184305 (2001).
- ⁸ K. Kudo, T. Noji, Y. Koike, T. Nishizaki, and N. Kobayashi, J. Phys. Soc. Jpn **70** 1448 (2001).
- ⁹ M. Hofmann, T. Lorenz, G. S. Uhrig, H. Kierspel, O. Zabara, A. Freimuth, H. Kageyama, and Y. Ueda, Phys. Rev. Lett. **87** 047202 (2001).
- ¹⁰ P. G. Klemens, Solid State Physics **7** 1 (1958), R. P. Tye ed., *Thermal conductivity*, Academic Press 1969, R. Berman, *Thermal Conduction in Solids*, Clarendon Pres, Oxford 1976
- ¹¹ J. Callaway, Phys. Rev. **113** 1046 (1959).
- ¹² Z. Rieder, J. L. Lebowitz, and E. Lieb, J. Math. Phys. **8**, 1073 (1967).
- ¹³ S. Lepri, R. Livi, and A. Politi, Phys. Rev. Lett. **78**, 1896 (1997), Europhys. Lett. **43**, 271 (1998), O. Narayan, and S. Ramaswamy, Phys. Rev. Lett. **89** 200601 (2002).
- ¹⁴ X. Zotos, J. Low Tem. Phys. **126** 1185 (2002).
- ¹⁵ K. Saito and S. Miyashita, J. Phys. Soc. Jpn, **71** 2485 (2002).
- ¹⁶ J.V. Alvarez and C. Gros, cond-mat/0201300/.
- ¹⁷ J.V. Alvarez and C. Gros, Phys. Rev. B **66**, 094403 (2002), Phys. Rev. Lett., **88**, 077203 (2002).
- ¹⁸ F. Heidrich-Meisner, A. Honecker, D. C. Cabra, and W. Brenig, Phys. Rev. B **66** 140406(R) (2002).
- ¹⁹ K. Saito, Europhys Lett. to be published.
- ²⁰ D. L. Huber, Prog. Theo. Phys. **39**, 1170 (1968), D. L. Huber and J. S. Semura, Phys. Rev. **182**, 602 (1969).
- ²¹ X. Zotos, F. Naef and P. Prelovsek, Phys. Rev. B **55**, 11029 (1997).
- ²² Exact treatment for all temperature regime may be possible using the argument of symmetries and the Mazur's inequality (P. Mazur, Physica **43** 533 (1969), see also the reference²¹). But here we confine ourselves to low energy regime which it is important in experiments.
- ²³ R. Kubo, M. Toda, and N. Hashitsume, *Statistical Physics II*, Springer-Verlag, New York, 1985.
- ²⁴ A. Klümper and K. Sakai, J. Phys. A **35** 2173 (2002).
- ²⁵ M. C. Cross and D. S. Fisher, Phys. Rev. B, **19** 402 (1979).
- ²⁶ T. Nakano and H. Fukuyama, J. Phys. Soc. Jpn, **49** 1679 (1980).
- ²⁷ A. Rosch and N. Andrei, Phys. Rev. Lett. **31** 1092 (2000).
- ²⁸ F. D. M. Haldane, Phys. Rev. B **25**, 4925 (1982)
- ²⁹ S. Sachdev, Quantum Phase Transition, Cambridge 1999.
- ³⁰ R. F. Dashen, B. Hasslacher, and A. Neveu, Phys. Rev. D **15** 4114 (1974).
- ³¹ Meisner et al.¹⁸ showed that for the general Tomonaga-Luttinger Hamiltonian, the Drude weight takes always $\frac{v\pi^2 T}{3}$ irrespective of the Tomonaga Luttinger parameter In the present calculation, this universal quantity can be confirmed.
- ³² S. Sachdev, and K. Damle, Phys. Rev. Lett. **78** 943 (1997).
- ³³ M. Garst, and A. Rosch, Europhys. Lett. **55** 66 (2001)
- ³⁴ E. Orignac, R. Chitra, and R. Citro, cond-mat/0211633/.